

[Web](#) · [Images](#) · [Groups](#) · [Directory](#) · [News](#)

Searched the web for **physostigmine amyloid**.

Results **111 - 120** of about **528**. Search took **0.23** seconds.

[PDF] Nicotinic Cholinergic Receptor Deficits in Alzheimer's Disease ...

File Format: PDF/Adobe Acrobat - [View as HTML](#)

... function produced by intracerebroventricular administration of β **amyloid** peptide in ... of a shorter acting, reversible AChE inhibitor (**physostigmine**) has been ...
www.mc.uky.edu/adreview/PDF/pauly.pdf - [Similar pages](#)

[[More results from www.mc.uky.edu](#)]

Research for Alzheimer's Disease - WrongDiagnosis.com

... of bases into the correct protein for beta-**amyloid**, causing a ... Another drug that inhibits acetylcholinesterase, called **physostigmine**, is helping researchers to ...

www.wrongdiagnosis.com/a/alzheimers_disease/research.htm - 25k - [Cached](#) - [Similar pages](#)

[PDF] Vol 172#2

File Format: PDF/Adobe Acrobat - [View as HTML](#)

... Neurobiologic features, such as the accumulation of **amyloid** and the reduction in ... inhibitors include first-generation compounds such as **physostigmine** and ta ...

mirecc.stanford.edu/mirecc/O'Hara%20AD%20Update.pdf - [Similar pages](#)

Affiliates - UW-Madison Institute on Aging

... Neuroendocrine responses to intravenous infusion of **physostigmine** in patients with ... effects on glucose metabolism, memory, and plasma **amyloid** precursor protein ...

aging.wisc.edu/research/affil.php?Ident=3 - 10k - [Cached](#) - [Similar pages](#)

NEJM -- Treatment of Alzheimer's Disease

... 2 Mutations in the gene for the **amyloid** precursor protein and the genes for presenilin 1 and 2 cause rare, dominantly inherited forms of the ... **Physostigmine**. ...

content.nejm.org/content/short/341/22/1670 - [Similar pages](#)

Herbal and Other Strategies for the Prevention and Treatment of ...

... The secondary events include the beta **amyloid** deposition, cytoskeletal and ... products area include rivastigmine (an analogue of **physostigmine**) '39 galanthamine ...

www.lightparty.com/Health/AlzheimerHerbal.html - 32k - [Cached](#) - [Similar pages](#)

Clinical Consult

... still unknown, researchers hypothesize that apolipoprotein E4 interacts with **amyloid-b** peptide ... The first AChE inhibitor studied was **physostigmine**, but its high ...

www.ascp.com/public/pubs/cc/1998/supp9a.shtml - 71k - [Cached](#) - [Similar pages](#)

[[More results from www.ascp.com](#)]

JPET -- Hirai et al. 280 (3): 1261

... B. and Nordberg, A.: Do tetrahydroaminoacridine (THA) and **physostigmine** restore acetylcholine ... Normal and abnormal biology of the beta-**amyloid** precursor protein ...

www.jpet.org/cgi/content/full/280/3/1261 - [Similar pages](#)

[[More results from www.jpet.org](#)]

Assigned Readings:

... Chromosome 21 - location of **amyloid** precursor protein gene ... 1) **Physostigmine** has been reported to be effective but had to be given iv Its duration of action was ...

www.uic.edu/classes/pmpr/pmpr652/Final/Koronkowski/alzheimer.html - 57k - [Cached](#) - [Similar pages](#)

[PDF] **B76236 Alzheimers 3/3**

File Format: PDF/Adobe Acrobat - [View as HTML](#)

... Reduction of β -**amyloid** peptide42 in the cerebrospinal fluid of ... and administration problems that bedevilled earlier treatments like **physostigmine** and tacrine. ...

www.alzheimer-insights.com/pdf/33.pdf - [Similar pages](#)

[[More results from www.alzheimer-insights.com](#)]




Result Page: [Previous](#) [2](#) [3](#) [4](#) [5](#) [6](#) [7](#) [8](#) [9](#) [10](#) [11](#) [12](#) [13](#) [14](#) [15](#) [16](#) [17](#) [18](#) [19](#) [20](#) [21](#) [Next](#)

[Search within results](#)

[Google Home](#) - [Advertise with Us](#) - [Business Solutions](#) - [Services & Tools](#) - [Jobs, Press, & Help](#)

©2004 Google



Entrez

PubMed

Nucleotide

Protein

Genome

Structure

PMC

Journals

B

Search

PubMed



for



Clear

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display

Citation



Show:

20



Sort



Send to

Text

Text Version

☐ 1: Brain Res Mol Brain Res. 1997 Jun;46(1-2):161-8.

Related Art

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

Pharmacological modulation of Alzheimer's beta-amyloid precursor protein levels in the CSF of rats with forebrain cholinergic system lesions.

Haroutunian V, Greig N, Pei XF, Utsuki T, Gluck R, Acevedo LD KL, Wallace WC.

Department of Psychiatry, Mount Sinai School of Medicine and B VA Medical Center, NY 10468, USA.

Abnormal deposition and accumulation of Alzheimer's amyloid beta protein (A beta) and degeneration of forebrain cholinergic neuro among the principal features of Alzheimer's disease. Studies in r model systems have shown that forebrain cholinergic deficits ar accompanied by induction of cortical beta-amyloid precursor pr (beta-APP) mRNAs and increased levels of secreted beta-APP : CSF. The studies reported here determined whether the CSF lev secreted beta-APP could be altered pharmacologically. In differ experiments, rats with lesions of the forebrain cholinergic syste received injections of vehicle, a muscarinic receptor antagonist scopolamine, or one of two cholinesterase inhibitors - diisoprop phosphorofluoridate (DFP) or phenserine. Scopolamine was administered to determine whether the levels of beta-APP in th could be increased by anticholinergic agents. The cholinesteras inhibitors were administered to determine whether the forebrain cholinergic system lesion-induced increases in CSF beta-APP c reduced by cholinergic augmentation. Scopolamine administratio a significant increase in the CSF levels of secreted beta-APP in lesioned rats. Phenserine, a novel, reversible acetyl-selective cholinesterase inhibitor, significantly decreased the levels of se beta-APP in the CSF of forebrain cholinergic system-lesioned r whereas DFP, a relatively non-specific cholinesterase inhibitor, to affect CSF levels of secreted beta-APP. These results sugge the levels of secreted beta-APP in the CSF can be pharmacolog modulated but that this modulation is dependent upon the status forebrain cholinergic system and the pharmacological properties drugs used to influence it.

MeSH Terms:

- Alzheimer Disease/metabolism
- Amyloid beta-Protein Precursor/cerebrospinal fluid*
- Amyloid beta-Protein Precursor/drug effects*
- Animals
- Cholinesterase Inhibitors/pharmacology*
- Isoflurophate/pharmacology
- Male
- Muscarinic Antagonists/pharmacology
- Physostigmine/analogs & derivatives
- Physostigmine/pharmacology
- Prosencephalon/drug effects*
- Prosencephalon/metabolism*
- Rats
- Rats, Sprague-Dawley
- Scopolamine/pharmacology
- Support, U.S. Gov't, Non-P.H.S.
- Support, U.S. Gov't, P.H.S.

Substances:

- Amyloid beta-Protein Precursor
- Cholinesterase Inhibitors
- Muscarinic Antagonists
- phenserine
- Scopolamine
- Isoflurophate
- Physostigmine

Grant Support:

- R01-AG10138/AG/NIA

PMID: 9191090 [PubMed - indexed for MEDLINE]

Display Citation ☒ Show: 20 ☐ Sort ☐ Send to Text

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

Dec 22 20